

# *Model Emerald Ash Borer Management Plan*

Jeffrey M. Hafner and J. Michael Orange

Updated September 2015

# **Model Emerald Ash Borer Management Plan**

Jeffrey M. Hafner and J. Michael Orange<sup>1</sup>

## **Acknowledgements**

The authors wish to thank the following people for their review and thoughtful comments on draft versions of this plan:

Professor Deborah G. McCullough, Ph.D., Department of Entomology, Michigan State University, [http://www.ent.msu.edu/directory/deborah\\_mccullough](http://www.ent.msu.edu/directory/deborah_mccullough)

Robert G. Haight, Ph.D., Research Forester, U.S. Forest Service, <http://www.nrs.fs.fed.us/people/haight>

Also, approximately 50 people who attended a two-hour work session during the Emerald Ash Borer Symposium held March 5 and 6, 2014 in Roseville, Minnesota vetted a draft of the plan.

Information provided here is for educational purposes only. References to commercial products or trade names do not imply endorsement by the authors or their institutions.

---

<sup>1</sup> Jeffrey M. Hafner is the Director of Municipal Consulting for Rainbow Treecare. J. Michael Orange is the Principal of ORANGE Environmental, LLC.

## Table of Contents

	<u>Page</u>
1. Introduction.....	1
2. The Emerald Ash Borer Infestation .....	3
2.1. Extent of the Infestation.....	3
2.2. Life Cycle of the Beetle and the EAB Death Curve .....	4
2.3. Experience of Other Cities.....	4
3. The Need for a Landscape-Based Management Strategy.....	6
4. The SLAM Study and Herd Immunity .....	7
5. The Model Emerald Ash Borer Management Plan.....	9
5.1. Goal 1—Accurate Tree Assessment and Record Keeping .....	9
5.2. Goal 2—Early Detection and Infestation Suppression .....	11
5.3. Goal 3—Postponement and Decrease of Peak Ash Mortality .....	14
5.4. Goal 4—Ash Tree Preservation While Minimizing Pesticide Use.....	15
5.5. Goal 5—Improvement of Tree Diversity and Expansion of the Tree Canopy .....	16
5.6. Goal 6—Minimization of Public Costs.....	17
5.7. Goal 7—Enlistment of the Private Owners .....	18
6. Pesticide Safety.....	19
7. Results of the Minnesota Ash Tree Preservation Analysis .....	19
8. Final Comment.....	20
References:.....	21
Figures:	
1. Net present value of surviving ash trees after five years for three different strategies.....	7
2. SLAM simulations .....	8
3. The Relationship between the EAB “Death Curve” and Management Options.....	20
Definitions.....	23

## 1. Introduction

The Minnesota Department of Natural Resources estimates that one in five (1.7 million) urban trees in the Twin Cities metro area is ash.<sup>2</sup> Without treatment, virtually all of these trees in the Twin Cities will be dead within ten years due to the invasive insect pest emerald ash borer (EAB), which has probably been in the area since 2006. Total Minnesota ash population estimates are close to one billion trees.

While the aesthetic value of trees is easily grasped, scientific studies have also quantified their benefits to the environment, the economy, and to human health. When the quantifiable benefits of trees are weighed against the costs (e.g. purchase, planting, pruning, and removal), the benefits outweigh the costs by a margin of about three to one. For example, an average mature ash will provide about \$170 in benefits annually.<sup>3</sup> The net cooling effect of a young, healthy tree is equivalent to 10 room-size air conditioners operating 20 hours a day.<sup>4</sup> One acre of urban forest absorbs 6 tons of carbon dioxide and emits 4 tons of oxygen annually.<sup>5</sup> Street trees even help extend the life of expensive asphalt by 40-60% by reducing daily heating and cooling of roads.<sup>6</sup>

Stormwater interception by trees reduces the peak-flow and flooding during intense storms, thereby reducing the amounts of pollutants that are washed into our rivers and lakes. Tree roots have a profound effect on the soil environment. They will direct 40-73% of assimilated carbon below ground,<sup>7</sup> and an average tree will intercept over 1,800 gallons of stormwater annually.

As regards human health, a recent analysis by the World Health Organization confirmed that air pollution is now the world's single largest environmental health risk.<sup>8</sup> In one study, stands of trees reduced particulates by 9-13%, and the amount of dust reaching the ground was 27-42% less under a stand of trees than in an open area. Another recent analysis, this one prepared by U.S. Forest Service scientists and collaborators, provides the first broad-scale estimate of how trees reduce air pollution, protect our health, and reduce health care costs. The article describing the analysis quoted Michael T. Rains, Director of the Forest Service's Northern Research Station and the Forest Products Laboratory: "With more than 80 percent of Americans living in urban area, this research underscores how truly essential urban forests are to people across the nation."<sup>9</sup>

---

<sup>2</sup> Source: "Rapid Assessment of Ash and Elm Resources in Minnesota Communities," Resource Assessment Unit, Forestry Division, Minnesota Dept. of Natural Resources, 1/12/07.

<sup>3</sup> National Tree Benefits Calculator, <http://www.treebenefits.com/calculator/>

<sup>4</sup> <http://www.arborday.org/trees/benefits.cfm>

<sup>5</sup> Ibid.

<sup>6</sup> Source: "City to Consider Special Funding for Trees," City of Madison Wisconsin, 7/31/14, <http://www.cityofmadison.com/news/city-to-consider-special-funding-for-trees>

<sup>7</sup> Source: [http://www.dailycamera.com/guest-opinions/ci\\_26131781/silent-environmental-devastation](http://www.dailycamera.com/guest-opinions/ci_26131781/silent-environmental-devastation)

<sup>8</sup> "7 million premature deaths annually linked to air pollution," World Health Organization press release, 3/25/14, [www.who.int/mediacentre/news/releases/2014/air-pollution/en](http://www.who.int/mediacentre/news/releases/2014/air-pollution/en)

<sup>9</sup> "Tree and forest effects on air quality and human health in the United States," Nowak, David, et al., *Environmental Pollution*, 7/25/14, <http://www.nrs.fs.fed.us/pubs/46102>

## Model Emerald Ash Borer Management Plan

The study estimated that in 2010, trees in the urban areas of Minnesota removed 4,600 tons of pollutants from the air and that this resulted in \$26.7 million in reduced health care costs.<sup>10</sup>

The above paragraph covers studies that quantify how trees benefit human health. Another study demonstrates how tree deaths from the infestation are associated with human deaths. The analysis by U.S. Forest Service scientists concluded that, “Poor air quality and stress are risk factors for [lower respiratory disease and cardiovascular disease], and trees can improve air quality and reduce stress. Results showed that the spread of EAB across 15 states was associated with an additional 15,000 deaths from cardiovascular disease and an additional 6,000 deaths from lower respiratory disease.”<sup>11</sup>

EAB will not only kill trees, it will severely affect the budgets of local governments; budgets that are already stressed by local economic bases that are still slowly recovering from the Great Recession. The negative impacts of EAB will extend beyond budget stress to include local governmental staff currently overwhelmed by constantly having to “do more with less.” The future holds even more peril considering that, over the course of this infestation, climate change effects and the costs to adapt to them will dwarf any previous challenges cities have faced. The future ability of Minnesota cities to effectively address the needs of their citizens while managing damage to infrastructure from invasive pests, an explosive increase in the number of hazard trees, and increased severe weather events is uncertain. The urban forest is one of the most valuable parts of the infrastructure of many Minnesota cities. For example, Minneapolis’ urban forest has an estimated value of \$756 million.<sup>12</sup> Most local governments that manage urban forests are woefully unprepared.

The EAB problem is very complex. Local physical and financial conditions will help define the most effective investment levels and timing regarding the following variables:

- Appropriate actions to reduce overall pest pressure
- Targeting for treatments those trees closest to the expected infestation wave front
- Subsidizing treatments for certain private trees
- Appropriate frequency of treatments and treatment doses
- Best practices for wood and debris handling issues

In the years soon after EAB was discovered in North American, most communities attempted to eliminate EAB through a single strategy—eliminating the food supply. It did not work, and recent research indicates the strategy is counterproductive.<sup>13</sup> Another strategy was to replace ash

---

<sup>10</sup> The health impacts and their monetary values are based on the changes in NO<sub>2</sub>, O<sub>3</sub>, PM<sub>2.5</sub> and SO<sub>2</sub> concentrations using information from the U.S. EPA Environmental Benefits Mapping and Analysis Program model (<http://www.epa.gov/air/benmap/>).

<sup>11</sup> “Exploring Connections Between Trees and Human Health,” *Science Findings*, Pacific Northwest Research Station, U.S. Forest Service, Jan./Feb. 2014, <http://www.fs.fed.us/pnw/sciencef/scifi158.pdf>

<sup>12</sup> Source: <http://www.americanforests.org/our-Programs/urbanforests/10-best-cities-for-urban-forests/10-best-cities-for-urban-forests-minneapolis/>

<sup>13</sup> At the Emerald Ash Borer Symposium held March 5 and 6, 2014 in Roseville, Minnesota, Dr. Daniel Herms, Professor and Chairperson of Entomology, Ohio State University, stated that it is a myth that preemptive removal of ash trees will slow the spread of the infestation. He argued that the removal of infested trees is effective but not the removal of healthy trees because the beetles will simply fly further to find host trees. He said that this would only spread the infestation faster. “We can’t cut

## Model Emerald Ash Borer Management Plan

trees with different species as fast as possible. However, a 2005 study of the urban forest in Minneapolis by the US Forest Service stated, “There is a delay of 30 years until the annual benefit of a replacement tree equals that of the ash tree removed because of EAB.”<sup>14</sup> The good news is that university scientists have developed and refined treatment protocols that can manage the infestation more effectively and at much lower cost when compared to past strategies that relied on preemptive removals or annual treatments for high-value trees.<sup>15</sup> We developed the Model Emerald Ash Borer Management Plan (Model EAB Management Plan) based on this new research.

The first part of this plan, Sections 2 through 4, provides a summary of the characteristics of the EAB infestation and two critical peer-reviewed scientific studies upon which this Model EAB Management Plan heavily relies. The first, called the Kovacs study, predicts that a regional or landscape-based management and funding strategy will be more cost effective in controlling an infestation than an inconsistent, city-by-city response. The second, known as the SLAM study, argues for an integrated pest management strategy that includes efforts to reduce pest populations, pesticide treatments, and other strategies to preserve valuable ash tree resources.

Section 5 describes the goals and the best practices for the Model EAB Management Plan and the science that forms the bases for them. Since there is concern for any plan that calls for the introduction of additional pesticides into the environment, Section 6 addresses pesticide safety. Appendix A includes a listing of terms used throughout this report. Appendices B and C summarize the components and the findings of a related analysis that estimates the costs, savings, and effectiveness of the integrated pest management strategy called for herein compared to a Base Case of not treating ash trees. Appendix D provides charts that illustrate these figures over the 20-year study period.

## 2. The Emerald Ash Borer Infestation and the Experience of Other Cities

**2.1. Extent of the Infestation:** The Emerald Ash Borer, *Agrilus Planipennis*, is an invasive beetle from Asia that was discovered in the United States during the summer of 2002 near Detroit, Michigan. It is the most destructive and economically costly forest insect ever to invade the U.S. By 2011, it was approximately 4 times as destructive nationally as the next two most costly pests, the Gypsy moth and the Hemlock adelgid.<sup>16</sup> The adult beetles ingest ash foliage but cause minimal damage. However, the larvae (during the beetle’s immature stage) feed on the inner bark of ash trees and disrupt the tree’s ability to transport water and nutrients. EAB may take years to build populations large enough to infest an entire tree, and low densities of EAB have little effect on the health of a tree

---

our way out of this.” Another presenter at the conference, Dr. John Bell, agreed and commented, “Don’t burn down the village to save it.”

<sup>14</sup> McPherson, E.G., Simpson, J.R., Peper, P.J., Maco, S., Gardner, S., Cozad, S., et al., 2005. City of Minneapolis, Minnesota Municipal Tree Resource Analysis. Center for Urban Forest Research, USDA Forest Service, Pacific Southwest Research Station. <http://www.treesearch.fs.fed.us/pubs/45984>

<sup>15</sup> For more information, refer to “Coalition for Urban Ash Tree Conservation,” [http://www.emeraldashborer.info/files/conserves\\_ash.pdf](http://www.emeraldashborer.info/files/conserves_ash.pdf)

<sup>16</sup> Aukema JE, Leung B, Kovacs K, Chivers C, Britton KO, et al. (2011), “Economic Impacts of Non-Native Forest Insects in the Continental United States,” *PLoS ONE*, 6(9): e24587. doi:10.1371/journal.pone.0024587, <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0024587>

(McCullough and Siegert 2007). However, once an ash tree is infested, it has almost zero chance of survival unless it is treated in time.

The current evidence from Michigan and Ohio shows that, depending on pest pressure, EAB takes five to ten years to infest and kill the majority of the ash trees in a city. Cities infested with this devastating pest have lost tens of millions of ash trees and endured billions of dollars of losses. As of January 2017, 29 states, including Minnesota and two Canadian Provinces, have discovered the EAB and enforced quarantines.

Many Minnesota cities have an abundance of ash trees on both private and public property, including boulevards. The complete loss of these trees due to EAB would have a devastating effect on home values, quality of life, and the environment. The time to act is now.

- 2.2. Life Cycle of the Beetle and the EAB Death Curve:** Recent research shows that the EAB beetle can have a one- or two-year life cycle. Adults begin emerging in mid-to-late May with peak emergence in late June. Females usually begin laying eggs about 2 weeks after emergence. Eggs hatch in 1-2 weeks, and the tiny larvae bore through the bark and into the cambium—the area between the bark and wood where nutrient levels are high. The larvae feed under the bark for several weeks, usually from late July or early August through October. Most EAB larvae overwinter in a small chamber in the outer bark or in the outer inch of wood. Pupation occurs in spring, and the new generation of adults will emerge in May or early June, to begin the cycle again.<sup>17</sup>

Cities that have been decimated by EAB have observed the EAB “death curve” where the rates at which infested trees die occur in two phases. EAB populations grow by a factor of 40 (or more) each year because the beetle has few natural predators and its host tree has limited natural defenses. However, healthy trees can tolerate an infestation for probably three to four years before they reach a tipping point that leads quickly to death. This results in a linear phase of the death curve where tree deaths are limited to about 1-5% a year. During the second phase of the death curve (the exponential phase), pest pressure builds, and tree deaths begin to parallel the exponential growth rate of beetle populations. Annual tree deaths can exceed 20%, and thousands of dead trees quickly overwhelm city crews, equipment needs, debris yards, and budgets. Most local governments that manage urban forests are woefully unprepared.

### **2.3. Experience of Other Cities**

We can learn from the experiences of other cities and states that have already been devastated by EAB. Westland, Michigan, one of the first communities hit by EAB, provides a specific example of the effects of the infestation. The removal of the 3,000 municipal ash trees killed by EAB “led to 33% increase in outdoor water consumption, which subsequently caused the regional water authority to levy a 10% surcharge on the

---

<sup>17</sup> Source: <http://www.emeraldashborer.info/faq.cfm#sthash.Bw4elMK7.dpbs>. See more at: <http://www.emeraldashborer.info/faq.cfm#sthash.Bw4elMK7.dpuf>

## Model Emerald Ash Borer Management Plan

city.”<sup>18</sup> Six counties in Ohio will spend \$1 million to clear 8,000 dead ash trees from the banks of a river because the dead trees were creating flooding problems.<sup>19</sup>

With EAB still sweeping through Illinois, Scott Schirmer, who is the Plant and Pesticide Specialist Supervisor and Emerald Ash Borer Program Manager at the Illinois Department of Agriculture’s Bureau of Environmental Programs, reflected about the state’s early experience: “We promoted local partnerships (cooperative efforts) for a long time as a way to deflect overall cost and hopefully spread benefits, but it never really panned out here. One of our major issues was the individuality of each municipality and arborist, as some were anti-pesticide and some were jumping on board. However, looking back, I truly believe that’s because the pesticides were still in their infancy and not ‘tried and true’ on a large scale at the time. Those communities that ‘had faith’ in the treatments are having great success today, whereas those that did not have lost large amounts of canopy. Communities on the leading edge have bought in a bit more, and are also benefitting from early, proactive treatment.”

Chicago is an example of an Illinois city that “had faith” in treatments. According to a press release from Mayor Rahm Emanuel, Chicago is using emamectin benzoate to inoculate its high-quality ash trees. “The City of Chicago is committed to the health of the more than 500,000 parkway trees that not only beautify our neighborhoods, but offer countless environmental benefits.”<sup>20</sup>

The City of Milwaukee began treating 33,000 of its larger ash trees in 2009 with emamectin benzoate as part of a plan to stage ash trees for their eventual removal while they transitioned to alternative species. However, residents have been so supportive of ash tree preservation that the city is considering altering the *Program* to long-term preservation and to begin protecting smaller ash trees as well.<sup>21</sup>

The experience of the City of Madison Wisconsin is telling as regards effects on the city budget, as reported in the *Wisconsin State Journal*. The newspaper quoted Mark Clear, a City Alderman, as stating, “One of the things really looming is the cost of the emerald ash borer. It’s one of the fastest growing areas of the budget, and it’s almost completely out of our control. I don’t think anyone wants to see the urban forest decimated.” The article went on to report that the city was removing approximately 8,500 ash trees and treating about 12,500 more. “The overall forestry budget is jumping from \$3.6 million in 2013 to a projected \$5.9 million in 2015.” To fund the increased costs, the city is

---

<sup>18</sup> Source: McCullough, Deborah G.; Mercader, Rodrigo J.; “Evaluation of potential strategies to SLow Ash Mortality (SLAM) caused by emerald ash borer (*Agrilus Planipennis*): SLAM in an urban forest,” *International Journal of Pest Management*, Vol. 58, No. 1, January–March 2012, 9–23.

<sup>19</sup> Source: <http://www.toledonewsnow.com/story/29364829/ash-trees-along-the-blanchard-river-to-be-removed>

<sup>20</sup> Chicago Office of Mayor Press Release, “City of Chicago begins second year of emerald ash borer inoculations: City to Treat 35,000 Ash Trees in 2014,” Source: <http://www.cityofchicago.org/content/dam/city/depts/mayor/Press%20Room/Press%20Releases/2014/May/5.19.14AshBorer.pdf>

<sup>21</sup> Source: “Managing EAB Risk with Ash Injections,” *Milwaukee Forestry*, Randy Krouse, Urban Forestry District Manager, City of Milwaukee, Wisconsin, [http://milwaukeeetrees.milwaukee.gov/pdf/Milwaukee\\_ManagEAB.pdf](http://milwaukeeetrees.milwaukee.gov/pdf/Milwaukee_ManagEAB.pdf)

considering levying a fee on property owners based on street frontage (which is even more regressive than property taxes).<sup>22</sup>

### **3. The Need for a Landscape-Based Management Strategy**

Trees are an integral part of the region's urban infrastructure and they should be viewed similarly to other components of our regional systems (land use, transportation, aviation, parks, and water resources).<sup>23</sup> The best approach to an EAB infestation is to fight it like a human health epidemic. Just as epidemiologists cannot fight a flu epidemic city by city, EAB cannot be efficiently fought city by city. While it is not necessary for 100% of the host population to be inoculated to control an epidemic, better results are achieved by inoculating a critical percentage of all hosts susceptible to the epidemic. That critical percentage is likely to be in the range of 20% of all ash trees in an area.<sup>24</sup> Since the beetle will kill virtually all untreated ash trees by the tenth year of an infestation, the percentage of treated trees relative to the total surviving ash population will eventually climb to 100%.

A recent scientific study, called the Kovacs Study, predicts that a regional or landscape-based management and funding strategy will more effectively control an infestation than an inconsistent, city-by-city response, or no response (Kovacs, Haight, Mercader, McCullough, 2013). The Kovacs study was based on the then-current EAB management plans and budgets for the ash trees located within a 10-kilometer radius of where EAB was first detected in the Twin Cities, which included St. Paul, Minneapolis, and 15 nearby cities. This analysis included multiple scenarios, three of which are critical here: The base case strategy assumed no infestation management plans or budgets and an unmitigated EAB population spread over a five-year period. The second strategy assumed individual cities managed the infestation independently with the known budgets at the time of the study and included treatments for 7% of the trees. The third strategy assumed the 17 cities pooled their resources to manage the infestation on a regional scale and included treatments for 24% of the trees.

As Figure 1 shows, the removal costs (\$5 million) in the base case far exceed the economic value of the remaining trees (not all of the ash trees will be dead after five years). The model returned a similar result if the cities acted independently and only treated 7% of the trees. However, when it was assumed in the third strategy that the cities pooled their resources and treated 24% of the trees in a manner that made sense for the region, the net present value of the surviving trees far exceeded the costs of removal. It is the comparison of the second and third strategies that makes this study unique. Even though both strategies assume the same amount is spent on management, the results are quite different. The report states that, "enabling municipalities to aggregate their

---

<sup>22</sup> Source: "Madison council members recommend special fee to pay for city's forestry program," *Wisconsin State Journal*, 7/31/14, [http://host.madison.com/wsj/news/local/govt-and-politics/madison-council-members-recommend-special-fee-to-pay-for-city/article\\_a66b8eb6-e485-5dd3-9b86-05743186d8a8.html](http://host.madison.com/wsj/news/local/govt-and-politics/madison-council-members-recommend-special-fee-to-pay-for-city/article_a66b8eb6-e485-5dd3-9b86-05743186d8a8.html)

<sup>23</sup> Since improved stormwater management constitutes one of the most important benefits from trees, the regional role in eliminating the combined sewer overflow problem offers an apt comparison. The SLAM study provided an example of the stormwater management benefit provided by trees: "In Westland, Michigan, one of the first communities adversely affected by *A. planipennis*, removal of the 3000 municipal ash trees killed by *A. planipennis* led to 33% increase in outdoor water consumption, which subsequently caused the regional water authority to levy a 10% surcharge on the city (T. Wilson, Westland Dept. of Public Works, pers. comm.)."

<sup>24</sup> Based on personal communication with Deborah G. McCullough, Ph.D., 1/17/14.

## Model Emerald Ash Borer Management Plan

budgets greatly improves total net benefits.... In addition, aggregate budget increases the percentage of healthy trees remaining in the final period by 18%, and the total net benefits more than double.”

The Kovacs report states that there is little active coordination among jurisdictions. We recommend that regional or state level public authorities formulate such a strategy so that it will be in place when the infestation begins to kill trees at an exponential rate.




Figure 1: Net present value of surviving ash trees after five years for three different strategies

### 4. The SLAM (SLow A.sh M.ortality) Study and Herd Immunity

The Model EAB Management Plan relies heavily on the scientific results from what is called the SLAM study (McCullough and Mercader 2012) and the study’s website of the same name (“SLAM Steps in Implementing a Strategy to SLow A.sh M.ortality”). As such, it is important to summarize the study before describing the components of the Model EAB Management Plan.

The SLAM study addressed the life cycle of the EAB beetle, pest pressure, the natural ability of healthy trees to tolerate an infestation, the limited effectiveness of sanitation to reduce the food source, the effectiveness pre-emptive removals and chemical treatments, the importance of early detection, and the relative costs of a variety of scenarios.


The SLAM website provides a succinct summary of the effort: “The rate at which ash tree mortality advances is related to EAB density.<sup>25</sup> Therefore, an over-riding theme within the

<sup>25</sup> From the SLAM study: “A low-density of *A. planipennis* [the EAB beetle] larvae generally has little effect on the overall health of the tree, in part because ash trees are highly sectorial (Tanis et al. forthcoming 2012) and relatively efficient at vertical translocation of nutrients and water. As larval densities build, however, more tissue is damaged, translocation is disrupted, canopies thin, branches die and eventually the tree succumbs.”

## Model Emerald Ash Borer Management Plan

SLAM approach is to reduce EAB numbers and the growth of EAB populations. ... A do-nothing or a regulation-only approach means that EAB populations will build and advance unchecked. Under that scenario, extensive local tree mortality is likely to occur much sooner than under a SLAM management scenario. ... The goal of this management strategy is to slow the local invasion process and allow land managers time to be proactive rather than simply reacting to overwhelming numbers of dead, often hazardous trees. ... Continued research and methods development will yield more options for EAB management and increase the effectiveness of existing technologies. Slowing the movement of EAB and the advance of ash mortality buys time for research and technology development” (“SLAM: SLow Ash Borer Mortality Pilot Project”)

The SLAM study included over 200 computer simulations based on field-derived data and derived a best-case scenario that was most effective at preserving ash trees at the lowest cost. This best-case scenario predicted that random treatment of 20% the population of ash trees annually should protect 99% of the trees after ten years. For comparison purposes, the study included a base case scenario that assumed no treatments. The costs of ash removal and replacement were “approximately fourfold higher than in any of the scenarios that included insecticide treatment. The dramatic difference in cumulative costs incurred, however, means that 20% of the ash trees could be treated for many years before treatment costs would approach removal and replacement costs” (McCullough and Mercader 2012).



**Figure 2:** Simulations represent scenarios in which systemic insecticides were applied annually to zero (No Treatment), 10%, or 20% of randomly selected trees. Source: “Evaluation of potential strategies to SLow Ash Mortality (SLAM) caused by emerald ash borer (*Agrilus planipennis*): SLAM in an urban forest,” Deborah G. McCullough and Rodrigo J. Mercader, *International Journal of Pest Management*, Vol. 58, No. 1, January–March 2012, 9–23.

While the SLAM study predicts success with limited treatments, its model is based on a known size and timing of the EAB infestation. Practitioners and policy makers in infested areas do not have access to such information and are forced to make a series of assumptions. The success of implemented plans will depend on the accuracy of these assumptions. The Model Emerald Ash Borer Management Plan was developed to provide tools to EAB managers that account for the unknown timeline and the uncertain growth rates of pest populations.

## Model Emerald Ash Borer Management Plan

Herd Immunity, also known as community immunity, is the public health phenomenon where protection from a disease for a critical percentage of the population allows protection for untreated individuals in the population. This principal occurs with a range of microscopic ‘bugs’ but the same concept applies to a larger bug—the Emerald Ash Borer. By treating a certain amount of the population of ash trees, there is a net benefit within the communities. The SLAM study calculates that 20% of the tree population is the amount needed to have a significant protective effect on the rest of the trees in the population.

### 5. The Model Emerald Ash Borer Management Plan

The purpose of the Model EAB Management Plan is to describe best practices for managing the infestation based on the most recent scientific findings. The Model EAB Management Plan also illustrates the cost advantages of using the full complement of integrated pest management strategies at the regional as well as local jurisdictional levels. The Plan is organized around seven goals, each with the accompanying best practices designed to implement them:

1. Maintain accurate assessments and records, and an updated city code.
2. Detect the infestation as early as possible and suppress the pest pressure.
3. Postpone and decrease peak ash mortality.
4. Preserve of the most valuable ash trees.
5. Expand the tree canopy and improve tree diversity.
6. Minimize public costs.
7. Enlist private ash tree owners to manage their trees consistent with the Ash Tree Preservation Plan.

- 5.1. Goal 1—Accurate Tree Assessment and Record Keeping, and Updating the City Code:** As with all infrastructure, maintenance is essential to maximize benefits, yet many cities lack the ability to track the maintenance and replacement needs of their urban forest. The Model EAB Management Plan links specific management strategies to categories of ash trees based on their condition, value, location, and ownership. Categories include woodlands, public trees and private trees, high- and low-quality trees, and high- and low-priority areas (refer to the Definitions section for explanations of each category). The i-Tree site is an excellent resource (<http://www.itreetools.org/index.php>).

#### Model EAB Management Plan Best Practices:

- **Public tree inventory:** Forest managers need a complete inventory of public trees that includes the size, species, condition, and location. This is essential for preserving this green resource and for identifying appropriate ash trees for use as detection and sink trees, for insecticide treatments, for preemptive removals, or for utilization for their wood value.
- **Private tree survey:** Management strategies of public trees can only affect a portion of total trees. Managers should obtain estimates of private trees based on surveys with a high degree of accuracy to know the extent of their influence.

## Model Emerald Ash Borer Management Plan

- **Ash tree assessment:** If budget or time constraints prevent a comprehensive tree inventory/survey, the first priority is to assess the ash trees located in high-priority areas, which are areas either within or within clear view from public lands and rights-of-way (boulevards, front yards of public and private property, and the mowed areas of public parks and open spaces).
- **Record keeping:** The implementation of an EAB management plan should include policies of regularly updating tree inventory database to reflect tree growth, removals, and replanting.
- **Updating the city code:** The infestation will likely require updates to the city code in the following ways:
  - **Nuisance language:** Few city codes will have up-to-date language regarding nuisances. Fortunately, the Minnesota Department of Agriculture and the League of Minnesota Cities have developed a model ordinance with specific language for “Control of Tree Infections and Infestations” and “Declared Tree Nuisances, Control Measures, and Control Areas.” The site includes specific definitions for the terms infestation and infection and are available for cities to copy and paste into their code.
  - **Upgrading landscape requirements in the zoning code:** At a time when it is important to maximize tree canopy as a major strategy to mitigate the effects of climate change, EAB will destroy thousands of trees. In order to take advantage of every opportunity to plant trees, the city can harness the power of the private sector through the development review process. The zoning code needs to incorporate all the best practices that maximize tree benefits. Numerous websites and other resources exist that offer state-of-the-art specifics to maximize tree canopy in development projects and parking lots.<sup>26</sup>
  - **On and off-site tree root protection zones:** Construction activities can damage trees in many ways. For example, soil compaction from vehicles and the storage of materials, and root severing, exposing, or burying can have long-term effects on a nearby tree’s roots such that it loses structural stability, becomes more susceptible to disease, and shorten its life. Construction impacts and the shortened life of a tree are often not obvious and inaction will continue to allow long-term damage to existing important trees.

Most codes will address the primary effects of construction activities (with a focus on erosion control) including regulations to protect important trees. Best practices require landscape plans identify trees both on-site and adjacent to the site that are classified as “Significant,” or “Desirable” based on generally acceptable arboriculture standards. Landscape plans should demonstrate how

---

<sup>26</sup> The Site Plan Review chapter of the Minneapolis Zoning Code (Chapter 530) offers an excellent example of how to maximize canopy cover through the development review process.

<http://library.municode.com/index.aspx?clientId=11490&stateId=23&stateName=minnesota&ds=site+plan+review>

## Model Emerald Ash Borer Management Plan

land-disturbing activities or permanent changes to the site would have no deleterious effect upon the tree root protection zones of important on-site or off-site trees.

- **Tree fund:** In the event site constraints prohibit the reasonable protection of on-site Significant and Desirable trees, the City could establish a tree fund and require developers to pay the equivalent costs for the City to buy, plant, and maintain trees on public or private property in the same watershed as the site.

These changes to the city code create roles for certified arborists in the development of landscape plans. This will slightly increase the costs of development but will significantly increase the quality of the landscape plans. This will make the job of city staff and decision makers easier because complete landscape plans will take the guesswork out of this aspect of the development approval process.

- **Alternative compliance:** It is very important to provide flexibility to accommodate unique site conditions and creative landscape designs. Cases will certainly arise where site constraints make it impossible or impractical to comply with all applicable requirements and still meet reasonable tree spacing needs (such as is needed for the health of the trees or to avoid underground or overhead utilities). Without the flexibility of an alternative compliance procedure, the increased rigor that would result from the other suggested text changes might discourage development or desirable landscape designs that meet the intent of the ordinances but not the letter of the law.

- 5.2. Goal 2—Early Infestation Detection and Suppression:** The SLAM study emphasizes the importance of early detection and actions to confine the infestation within the lessened pest pressure stage, which is the flat part of the death curve (refer to Section 2.2). This will reduce the vulnerability of the area to an overwhelming ash mortality event. Since appropriate control strategies vary according to the intensity of the infestation, it is important to detect the presence of the infestation as soon as possible and, once detected, periodically estimate the extent and the density of the EAB population.

Rather than remove low-value trees, they can serve a vital role as “trap trees.” Recent studies have shown that EAB beetles are attracted to stressed ash trees and tend to lay more eggs on stressed trees than on healthy trees.

### Model EAB Management Plan Best Practices:

- **EAB survey:** As soon as possible after initial detection, managers should conduct delimiting surveys organized in a grid pattern. Surveys should include visual inspections as well as some amount of destructive sampling (cutting and peeling of ash trees) to confirm the presence or absence of EAB.

## Model Emerald Ash Borer Management Plan

- **Tree girdling:** Research has shown that girdled<sup>27</sup> trees, especially those in sunny locations, are highly attractive to adult beetles in locations where EAB populations are relatively low. Girdled trees organized in a grid pattern are very effective for detection and assessment. The strategy of girdling trees can serve two purposes: 1) to assess beetle distribution, as well as larval density and development rates; and 2) to function as beetle population “sinks” to concentrate and eliminate adult beetles before they can disperse and reproduce.
- **Trap trees and population sinks:** Girdled trees are often referred to as “trap-trees,” a technique with a long history of use in forest pest management. The following are important considerations listed in the SLAM study:
  - If trees are girdled and remain standing for more than one year, they will serve as beetle magnets. Since girdled trees must be removed before the next generation of adults can emerge, a large component of future adults can be eliminated.
  - If tree cutting and removal is not a viable option, then creating *lethal* trap trees should be considered (refer to discussion below).
  - Girdled trees deployed in a systematic survey grid can concurrently serve as “sinks” for the subsequent generation of EAB.
  - Clustering three or four girdled trees creates a more powerful attraction for EAB adults than isolated single girdled trees in areas with low-density populations.
  - There is evidence to suggest that at very low EAB population levels, the location of sink trees can influence how beetles disperse. Sink trees will pull some beetles towards them as EAB adults respond to the presence of artificially damaged trees. Placing clusters of sink trees inside the core of an outbreak versus outside the outer edges could pull dispersing beetles away from the edges and potentially reduce spread rates.
  - Although all native ash trees will attract EAB adults, some species are more attractive than others. If different ash species are present, select by priority, from most to least preferred: (1) green ash, (2) black ash, (3) white ash, and (4) blue ash.
- **Timing for girdled trees:** Dates for girdling trap trees or setting traps and debarking trees or retrieving traps should be based on accumulated degree days for the local area since adults predictably fly at the same time each year. Girdled trees should be felled and debarked or destroyed in the fall, winter or early spring following their establishment to ensure that larvae die before completing development.

---

<sup>27</sup> Refer to the Definitions for a description of tree girdling.

## Model Emerald Ash Borer Management Plan

- **Removal of infested trees:** Infested trees must be removed or treated to ensure that developing EAB progeny are not allowed to emerge. Trees with canopy loss that exceeds 50% should be removed as soon as possible because dead ash trees generally deteriorate rapidly and many will become hazardous especially along streets, in yards, and along overhead power lines. This can entail chipping, grinding, debarking, burning or other methods.<sup>28</sup> Removing a few key infested trees, especially if they are large and heavily infested, could remove a locally significant number of EAB adults.<sup>29</sup> Since trees that have been dead for more than one year are unlikely to harbor EAB, their removal will not result in any reduction in the number of EAB.
- **Distant infestation:** The following guidelines apply if the closest known infestation is more than 15 miles away:
  - **Woodland detection trees:** Detection trees should be girdled in the early spring in accessible areas of woodlands, ideally in a grid pattern. Focus on areas closest to the expected wave front. Let trees die in place.
  - **High-priority area detection trees:** Same as above, but only girdle low-quality trees and remove them when they risk becoming hazard trees.
- **Proximate infestation:** The following guidelines apply if the infestation is within 15 miles or already within the jurisdiction:
  - **Woodland trap trees:** Girdle trap trees in the spring in accessible areas of woodlands, ideally in a grid pattern. Focus on areas closest to the wave front. Remove or process dead trees before adults can emerge in the spring.
  - **High-priority area trap trees:** Girdle low-quality trees in the spring and remove before adults can emerge in the spring.

**5.3. Goal 3—Postpone and Decrease Peak Ash Mortality:** Not all ash trees should be preserved. The Model EAB Management Plan incorporates an important strategy intended to reduce the overall intensity of the infestation, the pest pressure. Past strategies have included the removal of low-value ash trees to reduce the food supply. For ash trees in woodland areas and in low-priority areas such as backyards, a policy of “benign neglect” allows the EAB to kill the trees so the natural forest canopy can grow into the gaps. This can also reduce the current overabundance of ash in the urban forest. The problem with the “benign neglect” approach is that it allows EAB populations to increase exponentially wherever ash trees are left untreated. This increases overall pest pressure and hastens its spread.

---

<sup>28</sup> For information regarding firewood, refer to the following:

[http://www.nrs.fs.fed.us/disturbance/invasive\\_species/eab/control\\_management/survival\\_in\\_firewood/](http://www.nrs.fs.fed.us/disturbance/invasive_species/eab/control_management/survival_in_firewood/)

<sup>29</sup> “EAB infested trees can produce ca. 90-100 adult EAB per square meter (8-10 EAB per square ft.) of bark surface area. A single 20-inch diameter ash tree has the potential to produce approximately 3,600-4,000 beetles before it succumbs.” (“SLAM: SLow Ash Borer Mortality Pilot Project”).

## Model Emerald Ash Borer Management Plan

The Model EAB Management Plan includes the following best practices intended to reduce overall pest pressure, and to postpone and decrease peak ash mortality so forest managers can proactively manage the infestation rather than simply react to overwhelming numbers of dead, often hazardous trees.

### Model EAB Management Plan Best Practices:

- **Preemptive removals and ash utilization:** The first priority for low-quality trees in high-priority areas is for them to serve as detection/trap trees. Trees in low-priority areas can be preemptively removed for ash utilization and to reduce available phloem.<sup>30</sup> The removal of other trees can be staged as convenient over time.<sup>31</sup>

“Data from several sites have shown that in most areas, only a small proportion of ash trees are large (e.g. > 10 inches DBH),<sup>32</sup> while most ash trees are relatively small (< 4 inches DBH). Large ash trees can potentially produce hundreds to thousands of EAB adults but small ash trees produce relatively few, even when the small trees are abundant. Removing a few large trees can sometimes eliminate much of the available food for EAB larvae. Landowners may recognize some economic benefits by targeted harvests of large ash trees for lumber or firewood. ... Reducing ash phloem by itself is unlikely to slow spread. In some cases local EAB spread rates may increase because beetles are forced to fly further to locate a suitable host tree. An integrated approach that combines phloem reduction (e.g. removing selected trees) with insecticide treatments or girdling and sinks will be more effective than simply reducing ash phloem.” (“SLAM: SLOW Ash Borer Mortality Pilot Project”)

According to the SLAM study, ash trees are often common along road, railroad, utility, or trail right-of-ways, and that these types of corridors “enhance EAB dispersal and spread.” Therefore, they are excellent, accessible trees for preemptive removals and, if girdled, to serve as valuable sink trees.

- **Reducing pest pressure during moderate and peak periods:** As the infestation builds, it may be economically preferable to invest in reducing pest pressure near high-quality trees. Strategies include additional preemptive removals of low-quality trees (to reduce the food supply) and the use of trap trees. Lethal trap trees can be used by treating trap trees with emamectin benzoate a few weeks before girdling.

---

<sup>30</sup> From the SLAM study: “Sanitation to remove low vigour or hazardous ash trees can also be useful, especially in urban areas. Simply felling or removing ash trees to reduce the amount of ash phloem available for larvae, however, has the least effect on *A. planipennis* population growth and, if used exclusively, could increase spread rates.”

<sup>31</sup> Ash materials generated from tree removal should be disposed according to guidelines established by USDA Animal and Plant Health Inspection Service (APHIS)

([http://www.aphis.usda.gov/plant\\_health/plant\\_pest\\_info/emerald\\_ash\\_b/quarantine.shtml](http://www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b/quarantine.shtml)) and Minnesota Department of Agriculture. A properly managed community marshaling yard can enhance the disposal process.

<sup>32</sup> DBH refers to diameter at breast height i.e. 4.5 feet above the ground.

## Model Emerald Ash Borer Management Plan

However, the effectiveness of girdled trees to function as traps or sinks appears to diminish as EAB densities build in an area.<sup>33</sup>

SLAM study results indicate that achieving minimum overall treatment rates in an area (10-20% of all ash trees) can significantly reduce pest pressure. However, accomplishing even these seemingly low overall rates may require some degree of public investment in the management of private trees in the early years of the infestation before the beetles kill most of the untreated ash trees.

- **Strategies during periods of low pest pressure:** Strategies to reduce pest pressure, such as girdling and removing trap trees, can be expensive. Since ash trees can tolerate low levels of pest pressure, the best strategy is likely to invest only in treatments of high-quality trees closest to the likely wave front and inspections.
- **Encouraging natural enemies of EAB:** The SLAM study pointed out that treatments may increase the likelihood that beetle parasites and other natural enemies (e.g. beetle eating wasps and woodpeckers) can decrease beetle densities. Woodpeckers remain the most important natural enemy of EAB larvae, but woodpecker predation is not consistent. “Attracting woodpeckers into a local area and enhancing predation of EAB larvae could help to reduce EAB densities, it can also help in locating very lightly infested trees. Potential options for increasing woodpecker predation could include providing suet to retain woodpeckers in selected sites throughout the year. Classical biological control, which involves introducing a non-native natural enemy, may eventually be a part of SLAM efforts in some sites.” (“SLAM: SLOW Ash Borer Mortality Pilot Project”)

- 5.4. Goal 4—Preserve the Most Valuable Ash Trees:** The Model EAB Management Plan recommends reliance on trunk injection of emamectin benzoate (5 ml active ingredient per DBH) every 3 years to preserve important ash trees. Section 6 includes information regarding pesticide safety.

---

<sup>33</sup> For more detail, refer to McCullough, Deborah G.; Siegert, Nathan W.; *Using Girdled Trap Trees Effectively for Emerald Ash Borer Detection, Delimitation and Survey*, Dept. of Entomology and Dept. of Forestry, Michigan State University, July 2007. <http://www.emeraldashborer.info/files/handoutforpdf.pdf>

**Model EAB Management Plan Best Practices:**

- **Insecticide treatments:** Insecticide treatments should be used for the following public trees:
  - **Aggressive treatment protocol—Years 1 to 13:** Treat 100% of high-quality trees beginning with those closest to the infestation wave front, if known. Since trees can tolerate three or more years of low-to-moderate infestation, treat 1/3 of the trees each year to even out demands on crews, equipment, and budgets. Emamectin benzoate treatments are effective for three years or more (Herms, McCullough 2014).<sup>34</sup>
  - **Maintenance treatment protocol—Year 13 and beyond:** Inspect 100% of high-quality trees in Year 13. Treat (and track) those trees that show 30% or greater canopy decline thereafter. Implement SLAM Study practices by randomly selecting 20% of high quality trees for treatment in Year 13. Thereafter, treat 20% of randomly selected trees that had not been treated during the prior three years. Field research and the SLAM study confirm that treatments using emamectin benzoate will keep trees completely free of pests for the first two years after the injection, and that it takes three to four years after the start of an infestation for trees to decline to the degree they show at least 30% canopy loss and require removal.
- **Staging for removal and to serve as trap trees:** For urban forests where large numbers of ash trees are likely to need removal during the peak of the EAB death curve, forest managers may wish to treat trees so that they can survive long enough to be removed after the main wave of the infestation has passed. Continued inspection can determine when canopy loss exceeds 30%, after which they can be treated again to postpone removal, or girdled to serve as trap trees and then removed the following spring.

**5.5. Goal 5—Expand Tree Canopy and Improve Tree Diversity:** The tree diversity guideline known as the “10-20-30 rule” is an arboriculture guideline to reduce the risk of catastrophic loss due to pests like EAB. Since green ash, a single species, currently accounts for 15-20% of the urban forest (while the guideline suggests no more than 10%), the inevitable loss of virtually all untreated ash trees will eliminate this overabundance, and allow replacement trees to diversify forest and urban tree resources.

**Best Practice—Tree replacement:** Establish a policy that replaces trees in high-priority areas with at least a one-to-one ratio from a diversified list of eligible trees.

---

<sup>34</sup> A recent study examined the percentage of canopy decline in treated ash trees located in the midst of an EAB infestation (“Systemic Insecticide Technology for Tree Care,” by Dan Herms, Department of Entomology, Ohio State University, Ohio Agricultural Research and Development Center, Wooster, Ohio). The study found that trees treated with emamectin benzoate in 2006 (0.8 g ai / DBH) had only 5% canopy decline three years later in 2009.

- 5.6. Goal 6—Minimize Public Costs:** The Model EAB Management Plan is significantly less expensive and more effective than a remove-and-replace approach; and it preserves tree canopy and tree benefits. For the cost of removing and replacing two average 17” diameter trees, five mature trees can be preserved with treatments for over a decade.

**Model EAB Management Plan Best Practices:**

- **Budget balancing:** The following shows an approximate budgetary breakout by groups of best practices. It is intended to aid in the development of an EAB management plan for a local jurisdiction. During the implementation of the local jurisdiction’s plan, allocations should be expected to vary according to conditions on the ground. The percentage breakouts do not account for the costs of inventorying and estimating tree populations.
  - Cost of detection activities and the management of pest pressure: Approximately 15% of EAB management plan budget.
  - Cost of treatments, removals, and replacements: Approximately 80% of EAB management plan budget.
  - Cost of public outreach efforts: Approximately 5% of EAB management plan budget.
- **Record keeping:** A local government’s EAB management plan should include a tree inventory/survey. Proper record keeping over the course of the infestation will produce a trove of data that will be invaluable to the forest managers as well as other government officials and the scientific community as the knowledge base expands of how best to manage this infestation. It is an essential tool to battle the EAB infestation as well as future infestations and diseases. A wide variety of software programs exist for urban forest management, complete with standardized reports and the ability to customize them for EAB data recordation and evaluation.<sup>35</sup> The data needed to evaluate the EAB management program include the following:
  - **For each individual high-quality ash tree on public property:** Data should include geographic location, setting (e.g. boulevard, public yard or park, etc.), condition, size, management protocol (treatment in this case), treatment data (pesticide, treatment method, date of treatment, and dosage), inspection history, date of removal.
  - **Ash trees in low-priority public areas and low-quality ash trees:** Same as above.

---

<sup>35</sup> Davey Tree Co. has “TreeKeeper” software (c. \$2,000 per year): <http://www.davey.com/services/urban-forestry/urban-forestry-management-software/tree-keeper.aspx>

## Model Emerald Ash Borer Management Plan

- **Detection and trap trees:** Data should include geographic location, setting (e.g. boulevard, public yard or park, etc.), management protocol (e.g. girdling and removal).
- **Costs:** All program costs must be logged and tracked.
- **Public outreach history:** Records should include the program description, activity descriptions, and costs.
- **Program evaluation:** Careful record keeping will provide the data with which forest managers can compare the results on the ground with the predictions in the EAB management plan. For example, if the 3-year treatment protocol results in high-than-predicted canopy cover loss, the protocol should be changed to either every two years or the dosage per DBH increased. Similarly, some trees could serve as experiments in saving money by lengthening treatment periods or lowering dosages. A practice of early investments in detection and lowering pest pressure (e.g. through detection and trap trees) should be weighed against investing in treatments.

**5.7. Goal 7—Enlist Private Tree Owners:** A coordinated approach in urban areas will require a strong commitment to public outreach and education, especially in the years preceding the advent of the infestation and through its peak years.

### Model EAB Management Plan Best Practices:

- **Education and communication:** Governments should use all communication tools available to promulgate the goals and best practices in their EAB management plans, and to ensure that the owners of private ash trees manage their trees consistent with those plans. Educational and communication tools include the governmental website, newsletters, utility billings (for cities), and press releases. Community meetings are an excellent way to collaborate with those property owners most interested in preserving their ash trees.
- **Public subsidy for private trees:** Public forest managers may decide to subsidize treatments for certain ash trees on private property in order to help suppress pest pressure and to preserve certain trees. Only high-quality trees located in high-priority areas on streets where the loss of private ash trees would have a significant effect should be eligible for public subsidy. The tree inventory specific to the jurisdiction will provide the information needed to craft definitions for eligibility that will be most effective and enforceable. Budgetary constraints will determine the percent of the treatment costs to be subsidized. The subsidy should be contingent upon the property owner complying with the best practices described herein and should end after the third treatment when the peak of the infestation should have already occurred and the wave front will have moved on by the time the trees may need another treatment.

### 6. Pesticide Safety

We recognize the increasing and well-warranted concerns regarding the overreliance on pesticides. Neonicotinoids and their effects on pollinators, such as bees, and soil-applied products that have the potential to reach stormwater or ground water have all been highly publicized.

The pesticide recommended herein, emamectin benzoate (EB), is not a neonicotinoid and is injected into the trunks of the trees. Ash trees are wind pollinated, they are not a substantial nectar source for bees, and they flower early in the growing season and only for a limited number of days. “It is highly unlikely that bees would be exposed to systemic insecticides applied to ash.”<sup>36</sup> EB has a low toxicity rating for mammals, a low bioaccumulation potential within ecosystems, and is immobile in soil. This means that the insecticide will not build up levels within an ecosystem and will be minimally harmful to people and animals that might encounter tree debris.<sup>37</sup>

While there are valid concerns regarding the overuse of pesticides in our environment, those concerns should be aimed at reducing pesticide use where fewer benefits result. The environmental consequences of losing millions of ash trees are vastly greater than the minimal risk associated with inoculating high-quality ash trees to protect them from certain death. Marla Spivak, the Distinguished McKnight Professor in Entomology at the University of Minnesota, and an internationally recognized expert on bees, has said that the benefits of trunk-injected EB for ash trees outweigh the minimal potential harm to bees.<sup>38</sup> Dr. Deborah McCullough, a professor of entomology and forestry at Michigan State University, has stated, “There is no reason for a landscape ash tree to die from emerald ash borer anymore.”<sup>39</sup>

### 7. Results of the Minnesota Ash Tree Preservation Analysis

We analyzed the theoretical costs and benefits assuming the entire seven-county Twin Cities region participated in a landscape-based EAB management approach. Our Minnesota Ash Tree Preservation Analysis includes 100% of the approximately 450,500 public ash trees that are in the region’s urbanized areas.<sup>40</sup> The analysis compared two scenarios—a Base Case that assumed the removal of 100% of the ash trees and replacement of the 255,000 high-quality trees, and an integrated pest management approach called the Ash Tree Preservation Plan (ATP Plan). The

---

<sup>36</sup> “Frequently Asked Questions Regarding Potential Side Effects of Systemic Insecticides Used To Control Emerald Ash Borer,” <http://extension.entm.purdue.edu/eab/PDF/potentialSideEffectsofEABInsecticidesFAQ.pdf>

<sup>37</sup> “Emamectin benzoate is derived from a naturally occurring soil bacterium and has been registered for more than 10 years as a foliar spray to control pests in vegetable and cotton fields and parasitic sea lice in salmon aquaculture. Similar products are used in veterinary medicine as wormers for dogs, horses, and other animals.” “Insecticide concentrations that have been measured in treated trees are far below the levels known to be toxic to birds.... In Michigan and Ohio, where EAB has been established for several years, many ash trees have been treated with systemic insecticides. There have been no reported cases of woodpecker poisoning caused by insecticides applied for control of EAB.” (Hahn, Herms, McCullough, 2011)

<sup>38</sup> At the Minnesota Shade Tree Short Course held in Arden Hills, Minnesota, March 18 and 19, 2014, Dr. Marla Spivak said that the benefits of trunk-injected emamectin benzoate outweigh potential harm to bees.

<sup>39</sup> “Emerald ash borer treatments costing less, working better,” *Minneapolis StarTribune*, 8/8/13:

<http://www.startribune.com/local/south/218936301.html>

<sup>40</sup> Excluded were trees in non-urban areas and in woodlands, and private trees.

## Model Emerald Ash Borer Management Plan


ATP Plan scenario included pesticide treatments for the high-quality trees using trunk-injected emamectin benzoate. High-quality trees are public trees in good condition that are visible from active public areas (streets, front yards, and active areas of parks).<sup>41</sup>

Over the 20-year period of the infestation, the ATP Plan approach can preserve the high-quality trees at a total cumulative cost (\$145 million) that is over 40% lower than the Base Case cost (\$254 million) while preserving more than twice the tree canopy.<sup>42</sup> For every dollar of cost, the ATP Plan preserves approximately twice as much of the economic and environmental benefits from trees as the Base Case.

Using data from this analysis and from 17 Twin Cities suburbs, a typical city in the region is facing approximately \$1.5 million in EAB management costs over the next 10 years if it relies solely on a strategy of removal and replacement of city trees (the Base Case), or an \$800,000 savings if it employs the integrated pest management strategies described herein.

## 8. Final Comment

Acting individually, local governments can choose to manage the infestation or wait and let the beetles attack their already strained budgets. As the *Minnesota ATP Program* shows, the third and best choice is a holistic, landscape-based response that is centrally managed and that will minimize costs and maximize the value of the remaining urban forest. The *Minnesota ATP Program* will not only save money, it will reduce liabilities. The local government that delays action or relies on a removals-only approach will be overwhelmed with public hazard trees and probably the lawsuits that will follow. The time to act is now—before the infestation exponentially increases in population, and tree deaths escalate as seen in other cities. As the pest population increases and a greater number of trees die, the number of management options goes down (refer to Figure 3).



**Figure 3:** The Relationship between the EAB “Death Curve” and Management Options. Source: Davey Tree Company

## References

<sup>41</sup> Both scenarios assumed removal without replacement of approximately 195,500 low-priority trees. Low-priority trees are those in poor condition and those in areas that are not visible from publically accessible areas.

<sup>42</sup> The analysis uses trunk size as a surrogate for tree canopy measurements. Trunk size is measured as *diameter at breast height* (DBH).

## Model Emerald Ash Borer Management Plan

- “2010 Minnesota EAB Community Preparedness Manual,” Minnesota Department of Agriculture, Minnesota Department of Natural Resources, University of Minnesota, 5/18/10, <http://www.mda.state.mn.us/en/plants/pestmanagement/eab/eabmanual.aspx>.
- “EAB-Induced Ash Mortality in the Upper Huron River Watershed, SE Michigan,” OARDC, Ohio State University. [http://www.oardc.ohio-state.edu/hermslab/images/Herms\\_EAB\\_Management\\_12\\_Feb\\_2013.pdf](http://www.oardc.ohio-state.edu/hermslab/images/Herms_EAB_Management_12_Feb_2013.pdf). n.d.
- “Emerald Ash Borer.” <http://www.emeraldashborer.info>. n.d. December 20, 2013.
- “National Tree Benefit Calculator.” <http://www.treebenefits.com/calculator/treeinfor.cfm?zip=55118&city=SAINT%20PAUL&state=MN&climatezone=Midwest>. n.d.
- “Coalition for Urban Ash Tree Conservation,” [http://www.emeraldashborer.info/files/conserves\\_ash.pdf](http://www.emeraldashborer.info/files/conserves_ash.pdf)
- Cowles, Richard S., “The Facts About Systemic Insecticides and Their Impact on the Environment and Bee Pollinators,” Minnesota Turf and Grounds Foundation, Spring/Summer 2010. <http://www.ctpa.org/EAB%20Files/Clippings2010.pdf>.
- Doccola, Joseph J., Smitley, David R., Davis, Terrance W., Aiken, John J., and Wild, Peter M., “Tree Wound Responses Following Systemic Insecticide Trunk Injection Treatments in Green Ash (*Fraxinus pennsylvanica* Marsh) as Determined by Destructive Autopsy,” *Arboriculture and Urban Forestry*, 2011.37(1): 6-12.
- Doccola, Joseph J., Wild, Peter W., “Tree Injection as an Alternative Method of Insecticide Application,” Basic and Other Applications, Dr. Sonia Soloneski (Ed.), ISBN: 978-953-51-0007-2, InTech, 2012. <http://www.intechopen.com/books/insecticides-basic-and-other-applications/tree-injection-as-an-alternative-method-of-insecticide-application>.
- Frelich, Lee E. “Predicting Dimensional Relationships for Twin Cities Shade Trees.” Department of Forest Resources, University of Minnesota. [http://www.forestry.umn.edu/prod/groups/cfans/@pub/@cfans/@forestry/documents/asset/cfans\\_asset\\_249769.pdf](http://www.forestry.umn.edu/prod/groups/cfans/@pub/@cfans/@forestry/documents/asset/cfans_asset_249769.pdf). June 1992.
- Hahn, Jeffrey, Herms, Daniel A., McCullough, Deborah G., “Frequently Asked Questions Regarding Potential Side Effects of Systemic Insecticides Used To Control Emerald Ash Borer,” February 2011. [http://www.emeraldashborer.info/files/potential\\_side\\_effects\\_of\\_eab\\_insecticides\\_faq.pdf](http://www.emeraldashborer.info/files/potential_side_effects_of_eab_insecticides_faq.pdf)
- Herms, Daniel A., McCullough, Deborah G., “Emerald Ash Borer Invasion of North America: History, Biology, Ecology, Impacts, and Management,” *Annual Review of Entomology*, 2014. 59:13–30
- Herms, Daniel. “Systemic Insecticide Technology for Tree Care,” Department of Entomology, Ohio State University, Ohio Agricultural Research and Development Center, Wooster, Ohio. [http://www.oardc.ohio-state.edu/hermslab/images/Systemic\\_Insecticide\\_Technology\\_for\\_Tree\\_Care.pdf](http://www.oardc.ohio-state.edu/hermslab/images/Systemic_Insecticide_Technology_for_Tree_Care.pdf). n.d.
- Kovacs, Kent F., Haight, Robert G., McCullough, Deborah G., Mercader, Rodrigo J., Siegert, Nathan W., Liebhold, Siegert, Andrew M. Liebhold, “Cost of potential emerald ash borer damage in U.S. communities, 2009–2019,” *Ecological Economics* 69 (2010) 569–578.
- Kovacs, Kent F.; Haight, Robert G.; Mercader, Rodrigo J.; McCullough, Deborah G.; “A bioeconomic analysis of an emerald ash borer invasion of an urban forest with multiple jurisdictions.” *Resource Energy Econ.* (2013), <http://dx.doi.org/10.1016/j.reseneeco.2013.04.008>
- McCullough, Deborah G., Cappaert, D.L, Poland, T.M., Anulewicz, A.C., Lewis, P., and Molongoski, J., “Evaluation of Non-Invasive Trunk Sprays and Trunk-Injected Emamectin Benzoate,” meeting notes from the Emerald Ash Borer Research and Development Review Meeting, 2007.

## Model Emerald Ash Borer Management Plan

- McCullough, Deborah G.; Mercader, Rodrigo J.; “Evaluation of potential strategies to SLOW Ash Mortality (SLAM) caused by emerald ash borer (*Agrilus Planipennis*): SLAM in an urban forest,” *International Journal of Pest Management*, Vol. 58, No. 1, January–March 2012, 9–23.
- McCullough, Deborah G.; Siegert, Nathan W.; “Using Girdled Trap Trees Effectively for Emerald Ash Borer Detection, Delimitation and Survey,” Dept. of Entomology and Dept. of Forestry, Michigan State University, July 2007. <http://www.emeraldashborer.info/files/handoutforpdf.pdf>
- Rowntree, Rowan A. and Nowak, David J. “Quantifying the Role of Urban Forests in Removing Atmospheric Carbon Dioxide,” *Journal of Arboriculture* 17(10): October 1991. Accessed 3/22/13 at: [http://sfrc.ufl.edu/urbanforestry/Resources/PDF%20downloads/Rowntree\\_1991.pdf](http://sfrc.ufl.edu/urbanforestry/Resources/PDF%20downloads/Rowntree_1991.pdf).
- Sander, Heather A., Haight, Robert G., “Estimating the economic value of cultural ecosystem services in an urbanizing area using hedonic pricing,” *Journal of Environmental Management* 113 (2012) 194e205.
- “SLAM: SLOW Ash Borer Mortality Pilot Project,” <http://www.slameab.info/> December 22, 2013.

## Definitions

**Agrilus Planipennis:** An exotic beetle from Asia that was discovered in the United States during the summer of 2002 near Detroit, Michigan. Emerald ash borer (EAB) (Buprestidae: *Agrilus Planipennis* Fairmaire) is an exotic pest of ash (*Fraxinus* spp.). Feeding by the larval stage of this beetle occurs in tunnels called galleries that are excavated in the inner bark and phloem. The galleries created by feeding larvae affect the ability of trees to transport food and water. Low densities of EAB have little effect on the health of a tree. However, when EAB populations increase, the canopy declines, branches die and eventually the entire tree dies. Trees ranging from 1 inch to more than 60 inches in diameter have been killed by EAB.

**Cambium:** The cambium cell layer is the growing part of the trunk. It annually produces new bark and new wood in response to hormones that pass down through the phloem with food from the leaves. Refer to the following site for a description of tree structure: <http://www.arborday.org/treeGuide/anatomy.cfm>  
Condition rating: Condition rating refers to a system to rate the condition of trees from zero to four. A tree with a condition rating of four is the highest quality, zero is dead.

**EAB death curve:** The rates at which trees die occur in two phases: a linear phase (Years 1-4) and an exponential phase (Years 5-10).

**Emamectin benzoate:** A relatively new insecticide (sold as TREE-äge®) that can be used to protect valuable landscape ash trees from EAB. The product can be purchased and applied only by trained, certified arborists and landscapers. The product is applied as a trunk injection at the base of an ash tree. “A similar product has been used in salmon farming and in California for some fruit and vegetable crops. A related product is used in veterinary medicine.” Source: <http://www.emeraldashborer.info/treeage.cfm#sthash.MLsgMLtm.dpbs>.

**Girdling:** “Girdling, or removing a band of bark and phloem around the trunk of a tree, interrupts the ability of the tree to transport carbohydrates—the food needed by the tree. Girdled trees become increasingly stressed over the summer. As stress increases, the chemicals emitted from the foliage, bark or wood of the tree change. The wavelengths of light reflected by the leaves (hyperspectral reflectance) also differ between healthy and girdled trees. Beetles can apparently detect these changes and are often more attracted to the stressed trees than to surrounding or nearby ash trees. ... Recent studies by MTU and MSU scientists have shown that many ash trees can survive for at least two years after girdling. Trees can be girdled in the fall, winter or spring.” Source: McCullough, Deborah G.; Siegert, Nathan W.; *Using Girdled Trap Trees Effectively for Emerald Ash Borer Detection, Delimitation and Survey*, Dept. of Entomology and Dept. of Forestry, Michigan State University, July 2007. <http://www.emeraldashborer.info/files/handoutforpdf.pdf>

**Herd Immunity:** Also known as community immunity is the public health phenomenon where protection from a disease for a critical percentage of the population allows protection for untreated individuals in the population.

**High- and low-priority areas:** High-priority areas are areas either within or within clear view from public lands and rights-of-way. This includes boulevards, front yards of public and private property, and the mowed areas of public parks and open spaces. The term also applies to areas where it is especially important to preserve the environmental, economic, and aesthetic contributions from high-quality ash trees. Low-priority areas are areas that are not considered high-priority areas as regards the management of the EAB infestation.

**High-quality and low-quality trees:** High-quality trees refer to ash trees having a condition rating of 3 or greater located in high-priority areas. Low-quality trees are ash trees having a condition rating less than 3.

**Kovacs study and the landscape-based approach:** The response to the EAB infestation should be founded more on scientific conditions on the ground rather than constrained by governmental boundaries. A landscape-based focus can operate at all jurisdictional scales (state, region, city, neighborhood), and it incorporates both public and private trees (based on the Kovacs study<sup>43</sup>).

**Pest pressure:** The number of insects causing damage in one place at one point in time.<sup>44</sup>

- **Low pest pressure:** A condition in an area where beetle and larvae populations are relatively low such that the ash trees in an area can easily tolerate feeding levels and the associated phloem loss. A few trees may exhibit symptoms of infestation such as a few thinning branches but most trees appear healthy. Woodpecker holes may be present on the upper portion of the trunk or perhaps large branches on a few trees. EAB densities are low enough that most ash trees in the area will tolerate the associated phloem loss.
- **Moderate pest pressure:** A condition in an area where beetle and larvae populations are increasing. Symptoms of infestation will be apparent on several trees and a few trees may be seriously declining. Most ash trees in the area remain relatively healthy with less than 30% canopy dieback and/or thinning (transparency). If not treated with an effective pesticide, the pest pressure will continue to increase, leading to high mortality rates. If trees that remain relatively healthy (< 50% canopy dieback/thinning; little injury to the trunk) are treated with an effective product, trees should be able to recover (although dead branches will remain dead).
- **High pest pressure:** A condition in an area where beetle and larvae populations are at high densities. A few trees will be dead, a few others will be severely declining (and too injured to treat) and most others will exhibit canopy thinning and/or dieback. If not treated with an effective pesticide, most trees will die within 1-3 years. Trees that remain relatively healthy (<50% canopy decline) may recover, with the exception of the parts of the trees that suffered canopy loss.

**Phloem:** The food-conducting tissue of vascular plants, consisting of sieve tubes, fibers, parenchyma, and sclereids; also called bast.

**SLAM study:** The goal of the SLAM study was to “slow the onset and progression of ash mortality by slowing the growth of [beetle] populations.”<sup>45</sup> The study addresses the life cycle of the EAB beetle, pest pressure, the natural ability of healthy trees to tolerate an infestation, the effectiveness of preemptive removals and chemical treatments, the importance of early detection, and the relative costs of a variety of scenarios.

**Staging for removal:** During period of high pest pressure, there will be the greatest number of ash tree deaths. Urban forest managers may wish to treat trees so that they can survive long enough to be removed after the main wave of the infestation has passed.

**Tipping point:** the point where a tipping point when beetle larvae have consumed more than 60% of the phloem and the infested tree cannot recover. This leads inevitably to the death of the tree within a year or two.

**Trap trees as population sinks:** A strategy to moderate pest pressure is to girdle low-quality trees, already infested trees, and trees in low-priority areas such as woodlands to serve as trap trees. This is done in the fall in order to attract the adult beetles, which are especially drawn to stressed trees. The trees are felled the next spring and either processed or removed, thus killing concentrations of beetle larvae and the adults that are not yet mature enough to emerge.

---

<sup>43</sup> Kovacs, K.F., et al., “A bioeconomic analysis of an emerald ash borer invasion of an urban forest with multiple jurisdictions.” *Resource Energy Econ.* (2013), <http://dx.doi.org/10.1016/j.reseneeco.2013.04.008>

<sup>44</sup> Definitions are based on personal communication with Deborah G. McCullough, Ph.D., 1/17/14.

<sup>45</sup> “Evaluation of potential strategies to SLOW Ash Mortality (SLAM) caused by emerald ash borer (*Agrilus planipennis*): SLAM in an urban forest,” p. 11, op. cit.

## **Model Emerald Ash Borer Management Plan**

**Tree diversity:** The tree diversity guideline known as the “10-20-30 rule” is an arboriculture best practice designed to reduce the risk of catastrophic loss due to pests like EAB. The guideline recommends an urban forest be made up of no more than 10 percent of any one species, 20 percent of any one genus, and 30 percent of any one family. Since ash, a single species, currently accounts for about 20% of the urban forest, a significant reduction in numbers is appropriate per this guideline.

**Year X:** Refers to the years after the detection of the infestation within the jurisdiction.

**Woodland areas:** Areas where trees dominate and where development, mowed areas, and trails are absent or minimal. Topography and forest density makes protecting ash trees in woodlands difficult and uneconomical. However, to reduce overall citywide pest pressure, a minimum percentage of woodland trees should be girdled to act as traps and later removed and immediately processed to prevent adult beetles from dispersing. Beetles can emerge from ash logs or firewood for up to a year, and occasionally longer, after infested trees are felled. Woodland trees grow in close proximity to one another and compete for light. This competition reduces the canopy size of each tree, which makes the losses less significant to total canopy cover. Neighboring trees are positioned to quickly grow into the opened spaces created as ash trees die.